

DTIC FILE COPY

AFGL-TR-89-0004

AD-A205 189

Propagation and Attenuation of  
Lg Waves in South America

Ramon Roige Cabre  
Estela Ramos Minaya

Observatorio San Calixto  
Cas 5939  
La Paz, Bolivia

31 August 1988

Scientific Report No. 1

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

AIR FORCE GEOPHYSICS LABORATORY  
AIR FORCE SYSTEMS COMMAND  
UNITED STATES AIR FORCE  
HANSCom AIR FORCE BASE, MASSACHUSETTS 01731-5000

DTIC  
ELECTE  
21 FEB 1989  
A

89 2 21 077

Sponsored by:

DARPA Order No.:

Monitored by:

Contract No.:

Defense Advanced Research Projects Agency  
Nuclear Monitoring Research Office  
5299

Air Force Geophysics Laboratory  
Grant AFOSR-87-0311

The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the US Government.

"This technical report has been reviewed and is approved for publication"

  
JAMES F. LEWKOWICZ

Contract Manager

Solid Earth Geophysics Branch  
Earth Sciences Division

  
JAMES C. BATTIS

Acting Chief

Solid Earth Geophysics Branch  
Earth Sciences Division

FOR THE COMMANDER

  
DONALD H. ECKHARDT, Director  
Earth Sciences Division

This document has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA, Hanscom AFB, MA 01731-5000. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

# REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS													
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT  Approved for public release; distribution unlimited													
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		4. MONITORING ORGANIZATION REPORT NUMBER(S)  AFGL-TR-89-0004													
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. NAME OF MONITORING ORGANIZATION Air Force Geophysics Laboratory													
6a. NAME OF PERFORMING ORGANIZATION Observatorio San Calixto	6b. OFFICE SYMBOL (If applicable)	7a. ADDRESS (City, State, and ZIP Code)  Hanscom AFB Ma 01731-5000													
6c. ADDRESS (City, State, and ZIP Code)  Cas. 5939 La Paz, Bolivia		7b. ADDRESS (City, State, and ZIP Code)													
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Air Force Office of Scientific Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant AFOSR-87-0311													
8c. ADDRESS (City, State, and ZIP Code) Bolling Air Force Base Washington, DC 20332-6448		10. SOURCE OF FUNDING NUMBERS <table border="1"> <tr> <td>PROGRAM ELEMENT NO. 61101E</td> <td>PROJECT NO. 7A10</td> <td>TASK NO. DA</td> <td>WORK UNIT ACCESSION NO. DJ</td> </tr> </table>		PROGRAM ELEMENT NO. 61101E	PROJECT NO. 7A10	TASK NO. DA	WORK UNIT ACCESSION NO. DJ								
PROGRAM ELEMENT NO. 61101E	PROJECT NO. 7A10	TASK NO. DA	WORK UNIT ACCESSION NO. DJ												
11. TITLE (Include Security Classification) Propagation and attenuation of Lg waves in South America															
12. PERSONAL AUTHOR(S) CABRE <sup>~</sup> Ramón Roigé MINAYA Estela Ramos															
13a. TYPE OF REPORT Scientific Report #1	13b. TIME COVERED FROM 870801 TO 880731	14. DATE OF REPORT (Year, Month, Day) 1988 August 31	15. PAGE COUNT 36												
16. SUPPLEMENTARY NOTATION															
17. COSATI CODES <table border="1"> <tr> <th>FIELD</th> <th>GROUP</th> <th>SUB-GROUP</th> </tr> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td></td> </tr> </table>		FIELD	GROUP	SUB-GROUP										18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Lg, shield, cordillera, azimuth, path, particle motion, spectrum, SH, SV, Love, Rayleigh.	
FIELD	GROUP	SUB-GROUP													
19. ABSTRACT (Continue on reverse if necessary and identify by block number) <p>Lg study is introduced through a revision of investigations made since 1952 (when this phase was described for the first time). This study aims to know characteristics of Lg waves, as recorded in LPB station (WWSSN-type, 16.5°S, 68.1°W, 3292 m asl, with a gain of 50,000 or 25,000 time for SP and 1,500 times for LP).</p> <p>Epicenter lists were revised, sismograms inspected to annotate time, amplitude, period and group velocity of Lg phase. Several short-period records were digitized to draw particle motion and Fourier analysis.</p> <p>Earthquakes analysed were classified according to their characteristics into two groups coinciding with path type: a) across shields; b) across cordilleran zones; subdividing seismically active South America into the following origin zones: Brazil, Colombia and Venezuela, Chile, Argentina and Peru. --</p> <p style="text-align: right;">(continued on reverse)</p>															
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>													
22a. NAME OF RESPONSIBLE INDIVIDUAL James Lewkowicz		22b. TELEPHONE (Include Area Code) (617) 377-3028	22c. OFFICE SYMBOL AFGL/LWH												

UNCLASSIFIED

19. Continued

Period of maximal Lg waves changes between 0.5 and 2 seconds. Mean apparent group velocity is 3.6 Km/sec.


Particle motion is characterised by strong polarization horizontal, transverse to the direction of propagation.

For a shield path Lg amplitude is several times that of P; for a cordilleran path Lg is lesser than P, or also it does not appear at all.

UNCLASSIFIED

## Table of Contents

Introduction. Previous studies	p. 1
Objectives of the present study	3
LPB station	4
Method	4
Data used	5
Analysis	5
Fourier spectrum	10
Interpretation	10
Bibliography	12
Table I Lg velocity in different regions	15
Table II Earthquakes revised	16
Table III Lg mean apparent velocity	21
Fig. 1 Simplified geological map of South America	22
Fig. 2 Map of Lg paths across South America to LPB station	23
Fig. 3 Earthquakes recorded at LPB, travel across the shield	24
Fig. 4 Earthquakes, travel across the shield	25
Fig. 5 Earthquakes, travel across the Andes	26
Fig. 6 Particle motion: a) Brazil earthquake 11-20-80 b) Colombia 5-14-82	27
Fig. 7 Particle motion: a) Venezuela 7-17-79 b) Chile 6-28-72	28
Fig. 8 Particle motion: a) Argentina 12-05-77 b) Peru 6-27-83	29
Fig. 9 Lg spectra, earthquakes from Brazil and Colombia	30
Fig. 10 Lg spectra, Venezuela and Chile	31
Fig. 11 Lg spectra, Argentina and Peru	32



Grant N° AFOSR-87-0311

PROPAGATION AND ATTENUATION OF LG WAVES IN SOUTH AMERICA

Interim Technical Report

Ramón Cabré S.J., Principal Investigador

Estela Minaya, Associate Investigator

August 31, 1988

### Introduction. Previous Studies

Lg waves have been studied since 1952.

Press and Ewing (1952) describe for the first time Lg phase, introducing it as a shear wave train of period between 0.5 and 6 seconds, impulsive starting, with maximum amplitude among any other phases. They are recorded after distances of 6000 Km along continental structures. Group velocity amounts  $3.51 \pm 0.07$  Km/sec, with inverse dispersion, when distance source-station is more than  $20^\circ$ . Particle motion is predominantly transverse.

Lehmann (1953) remarked that vertical component of such waves is still significant.

Both (1954) analysed Lg phase, as recorded at Uppsala and Kiruna, noticing also a particle motion transverse. Both remarked two separate windows for group velocity:  $Lg_1$  of  $3.54 \pm 0.06$  Km/sec and  $Lg_2$  of  $3.37 \pm 0.04$  Km/sec.

Ewing et al. (1957), for Hungary and Crimea earthquakes, describe a motion predominantly horizontal transverse, accompanied by a lighter SV motion (so coinciding with Both (1954)).

Utzu (1960) finds a rather complicated particle motion.

Herrin and Richmond (1960) insist saying that Lg particle motion corresponds to transverse waves trending to some horizontal polarization.

The same again Waldner and Savarensky (1961).

On the other hand Koridalin (1961) shows samples of particle motion with weak horizontal polarization for  $Lg_1$  and strong polarization for  $Lg_2$ .

Pec (1962) focus particle motion as a strongly characteristic of SH.

For epicenters in Iran, Nuttli (1980) finds that the resulting wave amplitude is the double of vertical component.

Oliver and Ewing (1957) studied particle motion of  $M_2$ -mode, which is the first mode of shear waves, showing that it should be elliptic retrograde, as actually it was in a quake originated in the Arctic recorded at Palisades.

Group velocities, according to different authors, are listed in table I ; it appears that Lg group velocity is confined between a maximum 3.80 Km/sec and a minimum 3.19 Km/sec. Methods of calculating group velocity may change from one author to another and possibly method employed influences the results.

Also frequencies change for different authors (see table I ).

Several investigators find a relation of frequency to distance and to thickness of layer transmitting Lg.

Until now there is no definitive agreement about mechanism both of excitation and propagation of Lg; the main interpretations proposed are the following:

Press and Ewing (1952) introduce Lg as S waves with multiple reflexion within a granitic layer of the continental crust.

For Lehmann (1953) Lg is a superposition of different modes of Love waves.

Bath (1954, 1956, 1958) finds energy of  $Lg_1$  generally decreasing with focal depth increasing, energy of  $Lg_2$  is maximum for focal depth about 45 Km, what should be explained through the presence of several guide layers of different velocity.

Gutenberg (1955) suggested the presence of several low velocity guide layers, both in the crust and in the upper mantle.

Herrin and Richmond (1960) state that Lg is transmitted along the

upper crust, one to ten Km (or at most 14 Km).

Oliver and Ewing (1957, 1958), Yamaguchi (1961), Waldner and Savarensky (1961), Brune and Dorman (1963) all consider that a superposition of several higher modes of Love and Rayleigh waves explains frequency contents and group velocity.

Panza et al. (1972) consider that higher Rayleigh modes may be guided both along the crust and along mantle in a layer of a little lower velocity.

Knopoff et al. (1973) demonstrated that also Love waves may be transmitted as crustal and mantle channel waves. If no low velocity channel is in the mantle, upper modes of Love and Rayleigh waves should be responsible of Lg waves.

Panza and Calcagnile (1975) arrive to the same conclusion.

Ruzaikin et al. (1977) discard any usefulness of normal modes of surface waves to explain Lg; they suggest that lateral lithological changes along the path influence highly the characteristics of Lg.

Lg appears very relevant in some cases, so that magnitude of a seismic event and some underground characteristics of their path may be revealed; but in other cases Lg waves are not recorded at all; moreover some seismologists are claiming that under the same Lg name we are dealing with different kinds of seismic waves.

Studies of Lg waves across South America are really scarce: Cabré (1969-1971) emphasizes the influence of path on the recording amplitude.

Chinn et al. (1980) focus an especial case of Lg generation converted from Sn at the passage from an oceanic plate to the continental crust.

Raoof and Nuttli (1984-1985) have studied attenuation of Lg in South America.

#### Objectives of the present study

The main goals of the present study are:

a) To realize in what instances Lg is significant in the station LPB at La Paz. Bolivia, that is to say, from what seismic foci, across



what regions and with what amplitude Lg phase is clearly recorded.

b) To determine the nature of such waves.

A short synthesis of our results at this point will shed light to the whole report: Lg phase, as recorded in LPB station, is a group of short period waves, significantly large, of frequency 0.5 to 1.6 Hz, relevant for epicenters around the Brazilian or the Guiana Shields, with path mostly along the same shields, with velocity about 3.6 Km/sec, particle motion characteristic of the phase SH.

#### LPB Station

LPB is a WWSSN station installed in 1962 (replacing LPZ station located in a noisy site within La Paz town). Its coordinates are: -  
16°31'57.6"S; 68°05'54.1"W; 3292 m asl.

This station has operated quite regularly with a magnification of 50,000 times for short period 3 components until 1979, 25,000 times since then, and 1,500 times always for long period 3 components.

LPB is installed on a thick clay pan.

#### Method

During the elapsed first stage of research, the most important work consisted in selecting and observing records to distinguish characteristics of Lg waves, related to group velocity motion, energy spectrum and path from source to receiver.

1. Bulletin of the Observatorio San Calixto, Preliminary Determination of Epicenters of the U.S. Geological Survey and Bulletin of the International Seismological Centre were revised to list earthquakes originated in the region deemed to be able to produce Lg at the station LPB.

2. Corresponding seismograms were inspected to make a preliminary analysis of Lg phase.

3. The time of Lg phase was read and the characteristics of each case annotated, especially amplitude and period.

4. Apparent wave velocity epicenter-station was calculated in three ways:

- a) according to the arrival written, in some cases, in the bulletin;
- b) for the wave with maximum amplitude within Lg waves train;
- c) for the first arrival (be it emergent or impulsive), using as a criterion: the higher frequency appearing after longer waves (with some control of long-period seismograms) and the comparison of horizontal and vertical components.

5. Digitization of 3-component short-period seismograms for selected events at intervals approximately of 8 points per second, along about 20 seconds.

6. Drawing of particle motion.

7. Fourier analysis.

#### Data used

Short period records of 228 South American earthquakes between 1964 and 1986 were considered. Location, origin time and depth were accepted as published by the U.S. Geological Survey as Preliminary Determination of Epicenters (PDE) or in the Bulletin of the International Seismological Centre (ISC).

A few records, 14, were digitized in order to be Fourier analysed and particle motion reconstructed. They were originated:

Brazil 3	Chile 1
Colombia 3	Argentina 2
Venezuela 3	Peru 2

#### Analysis

Upon a quick inspection of seismograms a pathetic contrast of Lg recording appears for waves transmitted: a) across shields and b) across cordilleran zones (fig. 1) So, in the South American map the following zones were considered separately (see table II and fig. 2).

a) Seismic areas with path to La Paz across shield zones:

a-1. Brazil and Peru-Brazil Border:

14 Earthquakes were considered; the Nos. 1,2,6,9,10,14 have their wave path completely across the Brazilian Shield, N° 3 across the Guiana Massif, Wave arrival to LPB is from East and North-East. They have a Lg emergent but enough clear onset, unequivocally distinguishable from other previous phases.

Lg amplitude increases gradually until reaching a maximum (possibly interfering with Rg phase or surface waves, both of long and short period) and then amplitude decreases slowly along a longer duration - (fig. 3).

Maximum amplitude of Lg is three to four times that of P waves. The ratio Lg to S is still much larger, so that often S phase is not detected at all, meanwhile Lg is remarkable.

Lg waves are strongly polarized N-S, so that it is noticed by only an eye comparison of horizontal and vertical records.

Individual characteristics do not show any systematic relation to epicentral distance ( $10^{\circ}$  to  $33^{\circ}$ ), or to focal depth (5 to 65 Km), or to magnitude (mb 4.7 to 5.5); they do show a relation to azimuth epicenter-station ( $204^{\circ}$  to  $250^{\circ}$ ), without doubt because that yields a different wave path.

Group velocity for the largest waves within Lg was obtained, resulting a velocity higher than that obtained for other regions,  $3.6 \pm 0.2$  Km/sec (Table III).

That high velocity is an effect of the path, not of the origin; it agrees with the nature of rock across a crystalline, ancient basement (also P and S have a higher velocity than in other zones).

a-2. Colombia and Venezuela

The 94 earthquakes found in Colombia and Venezuela in the interval 1974 to 1984 are of special interest: (Table II) most of their path to LPB station is across the Guiana or Brazilian Shield, (fig. 2) and only a short way beneath the Andes.

Epicentral distance is between  $23^{\circ}$  and  $30^{\circ}$ .

Apparent velocity increases with depth (4 to 160 Km) as expected, but but

no other depth dependence is found for other characteristics.

The signature of those events is quite different from that appearing in Brazil Zone (fig. 3 and 4). Lg arrival is emergent, with 5 exceptions of impulsive arrival.

Waves are polarized EW.

Wave development may determine two typical envelopes; we call them A- or B- type. A-type shows a gradual increment of amplitude until a maximum where the phase Rg may interfere, and after that a slower decrement contrasts with previous increment. B-type does not reach a so large amplitude nor has a relevant maximum.

Generally Lg is larger than P and S (S is so attenuated that often it may not be distinguished on the record).

The mean apparent velocity is for both types 3.6 and 3.58 Km/sec (Table III).

The range of periods extends from 0.9 to 1.6 seconds.

Amplitude ratio Lg/P generally is larger than one (being Lg read at the vertical component and avoiding any confusion with Rg phase, what generally is much larger than Lg).

In some cases Lg is more attenuated than normal, but at the same time P is still more attenuated. In other cases Lg is so strong that it is seen on long-period records; it arrives before Rayleigh waves, several seconds after Love waves (that favors the identification of Lg with higher Love modes).

b) Seismic areas with cordilleran path to La Paz:

b-1 Central Chile

This area is the one with the most problematic Lg in LPB station.

Without doubt it is caused by its path along the Andes, of well known complexity both surficial and subcortical.

No characteristic or general behaviour may be established: Lg phase appears weak in some cases, but in many cases is not apparent at all, not being possible to state that it is not existent or it is so small that it disappears below the S-coda, since it should arrive less than 40 seconds after S and the threshold of readable Lg inscription is greater

than  $m_b = 5$  (in this part of seismogram SS, ScS, etc. are interfering ).

However Lg was clearly detected in 10 earthquakes (table II, fig. 2) with the following characteristics:

Epicentral distance is  $10^\circ$  through  $19^\circ$ ;

very emergent onset (fig. 5);

Lg may be larger or smaller than P;

waves polarized E-W;

Rg is not observed;

apparent velocity  $3.68 \pm 0.05$  Km/sec (Table III);

period between 1.0 and 2.0 sec;

some dispersion may be noticed in a few cases;

amplitude or duration of wave train does not increase significantly.

#### b-2 Central and South Argentina (Path to LPB parallel to the Andes)

This area is almost as complex as b-1; in the events recorded at LPB station (table II, fig. 2) Lg only hardly may be distinguished; its characteristics are different, but with more regularity, between an event and another.

94 Earthquakes from 1974 to 1984 were analysed, most of them originated in San Juan region (deep foci a priori were excluded, considering null the probability of any Lg phase). Often aftershocks followed the main shock after a few seconds, possibly masking Lg of the main event.

Epicentral distance is  $13^\circ$  through  $19^\circ$ ;

emergent onset; (fig. 5)

Lg is much smaller than P and S waves;

waves polarized E-W;

apparent velocity is 3.52 Km/sec (table III);

short period;

amplitude does not increase or decrease as much as in Brazilian events.

#### b-3 Peru and Peru-Ecuador Border

Among 14 events occurred between 1982 and 1986 (table II, fig. 2) Lg was recognized in only 7 (including one of epicentral distance less than  $10^\circ$ , considered minimum in other zones). In this case records from the station CCB sited at about 40 Km from LPB were also considered.

Epicentral distance  $8^\circ$  through  $18^\circ$  emergent onset (fig. 5);  
amplitude ratio Lg/P generally around one, but not systematic;  
mean apparent velocity  $3.6 \pm 0.3$  Km/sec;  
period 0.5 through 1.8 sec;  
amplitude changes are very small;  
no Rg is observed.

#### Generation and Propagation

Lg waves evidently are generated by a constructive interference of SH and SV, arriving to a low velocity layer (possibly the rock of upper most few kilometers); seismic waves are transmitted until recording station. Actual records are a result of both generation and propagation.

Indirect methods are necessary to know in what extent each process contributes to the final recording.

Lg waves period and velocity suggest a thin low velocity granitic layer (possibly 5 km thick), but the absence of that phase in some of the earthquakes of the region rather favors the interpretation that lateral changes of thickness, deep faults (such as Oca fault in N Colombia - Kellog et al. 1982), gradual metamorphism, sedimentary basins are responsible of changing amplitude for seisms of the same region.

#### Particle Motion

Lg phase is strongly polarized horizontally for all the paths considered.

So, particle motion is characteristic of all Lg phases, independently of the origin region (figs. 6, 7, 8).

during several seconds of initial Lg waves SH predominates on SV; later on a new arrival, now both at the same time, of SH and SV; after that SV component is stronger than SH.

We shall quote Nuttli (1961) and Nuttli et al. (1962): "According to this relation the surface motion produced by an incident S wave may be considered in two separate cases: 1. Angle of incidence less than  $\sin^{-1}(b_0/a_0)$ , where  $a_0$  and  $b_0$  are the efficient P and S wave velocities at the earth's surface. For this case

there is generated both a reflected P and a reflected S, and the surface particle motion is linear.

2. Angle of incidence larger than  $\sin^{-1} (b_0/a_0)$ . For this case there is total reflexion of the incident S wave. In general the three components of the earth's surface, namely the SH and the horizontal and vertical components of SV, are out of phase with respect to one another. The surface particle motion is non linear".

#### Fourier Spectrum

Spectrum contents (figs. 9, 10, 11) is analysed for all 14 cases as a whole; amplitude logarithm is plotted versus frequency. A maximal pick appears in general between 0.77 and 0.45 Hz; that value does not agree with the first appearance of the record, being the difference originated in the instrumental response curve and in the filtering used (it amounts about ten percent, and may not influence significantly our interpretation in the present research period).

#### Interpretation

In general, we have realized for South America what has been said by different authors for other regions:

1. Earthquakes originated well inside an oceanic crust do not produce any Lg phase.
2. Cordilleran structure attenuates so much Lg waves that for a long path across it Lg either is very small or not existent.
3. For surface foci small variations in depth influence strongly Lg generation.
4. According to the precedent statement and to the wave velocity, Lg is a train of guided waves in a crustal layer.
5. Particle motion maintains some relation both to the angle of incidence and P and S velocities

#### Specifically we find:

- a) Earthquakes originated North and East of the Brazilian Shield have a prominent phase Lg in the LPB station. Those zones mean politically Northern Ecuador, Colombia, Venezuela, Guianas, Eastern Brazil.
- b) For earthquakes originated in Western Brazil (Brazil-Peru border) Lg is really small, or not existent at all.

c) Colombia earthquakes have large Lg also for intermediate depth foci; in other zones Lg amplitude decreases rapidly for increasing depth.

d) Argentinian or Chilean earthquakes, having a path to La Paz mostly beneath the cordillera, do produce really small waves Lg.

e) Lg appears well transmitted across metamorphic and igneous rocks not strictly granitic.

f) Particle motion is characteristic of all phases Lg, independently of the region considered. It is identical to that of SH and well distinguishable since SH arrives before SV and is better developed.

g) In the case of large Lg waves it seems that they increase by interference of S waves propagation forward and those reflected back in some discontinuity.

h) For the Venezuela area it is impossible to separate SH and SV; it is not clear if this impossibility arises from the superposition of SH and SV, or possibly from the superposition of Pn, Sn and Love codas.

i) Being LPB station within a cordilleran zone, though close to its border, the question how Lg waves are recorded so large deserves a longer discussion.

We did not find a convincing explanation of what happens with a train of waves arriving to a medium of high attenuation, but the answer to this question is well known in a similar case: in a local quake the more attenuant is the medium the larger are waves in it (a pathetic example is the town of Mexico destructed by a quake distant about 400 Km, but only destructed in the part where soft underground has a high attenuation). That should mean for our problem that energy conveyed by Lg waves is transformed into shaking amplitude until it is completely dissipated, say not apparent at all after about 200 Km; that contrasts with previous transmission along a regular guide layer, with very little attenuation.



# B I B L I O G R A P H Y

- Båth, M. (1954), The elastic waves Lg and Rg along Euroasiatic paths, Arkiv. Geophys., 2, 295-324.
- Båth, M. (1956), Some consequences of the existence of low velocity layers, Ann. Geofis., 9, 411-450.
- Båth, M. (1958), Channel waves, J. Geophys. Res., 63, 583-587.
- Båth, M. (1959), Seismic channel waves. New observations and discussions, Beitrage zur Geophysik, 68, 360-376.
- Bollinger, G.A. (1979), Attenuation of the Lg phase and the determination of  $m_b$  in the Southeastern United States, Bull. Seism. Soc. Am., 69, 45-63.
- Bolt, B. (1957), Velocity of seismic waves Lg and Rg across Australia, Nature, 180, 495.
- Bolt, B., H.A. Doyle, and D.F. Sutton (1958), Seismic observations from the 1956 atomic explosions in Australia, Geophys. J.R. Astr. Soc., 1, 135-145.
- Brune, J. and J. Dorman (1963), Seismic waves and earth Structure in the Canadian Shield, Bull. Seism. Soc. Am., 53, 167-210.
- Cabr , R.S.J. (1969), Ondas Lg Registradas en La Paz, Bolivia, Tesis de Doctorado, Universidad de Barcelona, Espa a.
- Cabr , R.S.J. (1971), Ondas Lg Registradas en La Paz, Bolivia. Geof sica Panamericana. 1, 71-95.
- Chinn, D.S., B.L. Isacks and M. Barazangi (1980), Highfrequency seismic wave propagation in western South America along the continental margin, in the Nazca Plate, and across the Altiplano, Geophys. J.R. Astr. Soc., 60, 209-244.
- Ewing, M., W.S. Jardetzky, and F. Press (1957), Elastic Waves in Layered Media, MC Graw-Hill, 219-222.
- Gane, P.G., A.R. Atkins, J.P.F. Sellschop and P. Seligmann (1956), Crustal Structure in the Transvaal, Bull. Seism. Soc. Am., 46, 293-316.
- Gamber, F. and P.W. Pomeroy (1970), Seismic wave velocities and earth structure on the African Continent, Bull. Seism. Soc. Am., 60 651-668.
- Gutenberg, B. (1955), Channel waves in the Earth's crust, Geophys., 20, 283-294.
- Herrin, E. and P.D. Milton (1960), The velocity of Lg in the Southwestern United States and Mexico, Bull. Seism. Soc. Am., 50 35-44.
- Herrin, E. and J. Richmond (1960), On the propagation of the Lg phase, Bull. Seism. Soc. Am., 50, 197-210.
- Herrin, E. (1960) On  $\bar{P}$  and Lg, J. Geophys. Res., 66, 334-335.
- Hodgson, J.H. (1953), A Seismic Survey in the Canadian Shield II: Refraction Studies based on timed blasts, Publ. Dominion Observatory, Ottawa, 16, 167-181.

- Horner, R.B., A.E. Stevens, and H.S. Hasegawa (1973), The Bengough, Saskatchewan earthquake of July 26, 1972, *Can. J. Earth Sci.*, 10, 1805-1821.
- Kellogg, I.N. and W.E. Bonini (1982), Subduction of the caribbean plate and basement Uplifts in the overriding South American Plate, *Tectonics*, 1 N° 3, 251-276.
- Knopoff, L., F. Schwab and E. Kausel (1973), Interpretation of Lg, *Geophys. J.R. Astr. Soc.*, 33 389-404
- Knopoff, L., F. Schwab, K. Nakanishi, and F. Chang (1974), Evaluation of Lg as discriminant among different continental crustal structures, *Geophys. J.R. Astr. So.*, 39, 41-70.
- Koridalin, E.A. (1961) Some characteristics of Lg and Rg waves and regional features in their propagation, *Izv. Geophys. scr.*, 1114-1121.
- Lehamann, I. (1953). on the Short-period surface wave "Lg" and crustal structure, *Bull. d' Information, U.G.C.I.*, 2, 248-251.
- McEvelly, T.V. (1964), Central U.S. Crust - upper mantle structure from Love and Rayleigh wave phase velocity inversion, *Bull. Seism. Soc. Am.*, 1997 - 2015.
- Nuttli, O.W. (1956), Tentative velocities of Seismic Crustal waves in the Central United States, *Earthquake notes* 27, 41-44.
- Nuttli, O.W. (1961), The effect of the Earth's Surface on the S wave Particle motion, *Bull. Seism. Soc. Am.*, 51, 237-246.
- Nuttli, O.W. and J.D. Whitmore (1962), On the determination of the polarization angle of the S wave, *Bull. Seism. Soc. Am.*, 52, 95 - 107.
- Nuttli, O.W. (1980), The excitation and attenuation of seismic crustal phases in Iran, *Bull. Seism. Soc. Am.*, 70, 469-485.
- Oliver, J.E., M. Ewing and F. Press (1955), Crustal structure of the Arctic regions from Lg phase, *Geol. Soc. Am.*, 66 1063-1074.
- Oliver, J. and M. Ewing (1957), Higher modes of continental Rayleigh waves, *Bull. Seism. Soc. Am.*, 47, 187-204.
- Oliver, J. and M. Ewing (1958), Normal modes of continental surface waves, *Bull. Seism. Soc. Am.*, 48, 33-49.
- Panza, G.F., F.A. Schwab, and L. Knopoff (1972), Channel and crustal Rayleigh waves, *Geophys. J.R. Astr. Soc.*, 30, 273-280.
- Panza, G.F. and G. Calcagnile (1975), Lg, Li and Rg from Rayleigh modes, *Geophys. J. Roy. Astr. Soc.*, 40, 475.
- Payo, G. (1960), Estudio sobre las ondas superficiales Lg, Rg y Li en los registros del Observatorio de Toledo, *Rev. Geofísica*, 5, 227.
- Pec, K (1961), Phase Lg et Rg Observées a Praha, *Ceskoslovenská Akad. Ved. Studia Geophys. et Geod.*, 5, 219-226.
- Pomeroy, P.W. and T. Novak (1978), An investigation of Seismic wave propagation in western USSR, *Semi-Annual Technical Report N° 2*,

- Rondout Associates, Inc. Stone Ridge, NY.
- Press, F. and M. Ewing (1952), Two Slow surface waves across North America, Bull. Seism. Soc. Am., 42, 219-228.
- Press, F. (1956), Velocity of Lg waves in California, Trans. Am. Geophys Union, 37, 615-618.
- Raouf, M.M., O.W. Nuttli (1984-85), Attenuation of high-frequency Earthquake waves in South America, PAGEOPH, 22, 619-644.
- Rizikova, S. (1966), The velocities of the Seismic waves Li, Lg and Rg across the Aegean Sea, Bull. del 1' Inst de Géophys. Acad. Bulg. Sci, 2.
- Rizikova, S. (1968), Period dependencies of the Li, Lg<sub>1</sub>, Lg<sub>2</sub> and Rg, Bull. de 1' Inst de Géophys. Acad. Bulg. Sc., 12, 113-124.
- Ruzaikin, A.I., I.L. Nersesov, V.I. Khalturin and P. Molnar (1977), Propagation of Lg and Lateral variations in crustal structures in Asia. J. Geophys. Res., 82, 307-316.
- Savarensky, E.F. and N.G. Waldner (1960), Lg and Rg waves from earthquakes in the Black Sea basin and certain consideration on their nature, Symp. "Seismic Studies in the period of the IGY", 4 Acad. Sc., USSR Press.
- Stauder, W. and G. Bollinger (1963), Pn velocity and other seismic studies from the data of recent Southeast Missouri earthquakes, Bull. Seism. Soc. Am., 53, 661-679.
- Utzu, T. (1960), Observations on the Lg waves and crustal structure in the vicinity of Japan, Geophys. Mag., 29, N° 4.
- Waldner, N.G. and E.F. Savarensky (1961), On the nature of the Lg<sub>1</sub> wave and of its propagation in Northeast Asia, Izv. Geophys. Ser., 3-24.
- Yamaguchi, R (1961), Surface waves and layered structures, Part I, Bull. Earthq. R. Institute, 39, 653-668.
- Willmore. P., A.L. Hales and P.G. Game (1952), A Seismic investigation of structure in the western Transvaal, Bull. Seism. Soc. Am., 42, 53-80.

Table I Lg VELOCITY IN DIFFERENT REGIONS

Region	Velocity (Km/sec)	Period (sec)	References
Africa	3.48 to 3.60		Gumper y Pomeroy (1970)
S. Africa Transvaal	3.68		Willmore et al (1952)
" " "	3.66		Gane et al (1956)
Australia	3.50	3 to 6	Bolt (1957)
"	$3.44 \pm 0.04$		Bolt et al (1958)
Eurasia	$3.54 \pm 0.06$	Lg <sub>1</sub>	Bäth (1954)
"	$3.37 \pm 0.04$	Lg <sub>2</sub>	" "
N America	$3.51 \pm 0.07$	5 to 6	Press Ewing (1952)
Eastern N America	3.57	0.5 to 1 for 19.6°	Lehmann (1953)
		5 for 35°	" "
Canadian Shield	3.54		Hodgson (1953)
" "	3.60 to 3.70		Brune and Dorman (1963)
" "	3.56		Horner et al (1973)
California	$3.54 \pm 0.02$		Press (1956)
Sierra Nevada	$3.53 \pm 0.02$		" "
Central Valley and Coastal Ranges	$3.55 \pm 0.03$		" "
Central U.S.	3.49		Nuttli (1956)
" "	3.49 to 3.80		McEvelly (1964)
" "	3.03 to 3.39		Pomeroy and Novak (1978)
" "	2.18 to 3.72		" " "
Eastern U.S.	3.19 to 3.35		" " "
" "	3.04 to 3.80		" " "
Central and SE U.S.	$3.65 \pm 0.04$		Stauder and Bollinger (1963)
SE U.S.	$3.50 \pm 0.13$	$0.7 \pm 0.1$	Bollinger (1979)
" "	$3.52 \pm 0.10$	$0.8 \pm 0.1$	" "
Hungary and Crimea	3.16, 3.51, 3.23		Bäth (1959)
Coast of Mexico	3.47 to 3.54	1 to 7	Herrin and Richmond (1961)
Kamchatka Japan and Kuril Is.	3.50 Lg		Waldner and Savarenstky (1961)
" "	3.29 Lg <sub>2</sub>		" " " "
S China	3.53 Lg <sub>1</sub>	6 to 10	" " " "
" "	3.31 Lg <sub>2</sub>		" " " "
Sofia, Bucarest,			
" Istanbul	3.57 Lg <sub>1</sub>		Rizikova (1966-68)
" "	3.32 Lg <sub>2</sub>		" "
Europe	3.61		Payo Subiza (1960)
"	3.36		" " "
South America	3.8 to 4.4 early Lg		Chinn et al. (1980)
" "	3.6 Lg		" " "
" "	3.5 Late Lg		" " "
Black Sea Basin	3.60 Lg <sub>1</sub>	15°	Gutenberg (1955)
" " "	3.55 Lg <sub>1</sub>		Savarensky et al (1960)
Vicinity of Japan	3.51 Lg <sub>1</sub>		Utzu (1960)
Eurasia and Arctic	$3.54 \pm 0.03$	1 to 9	Koridalin (1960)
Artic	3.48		Oliver et al (1955)
South West U.S. and Mexico	$3.47 \pm 0.01$		Herrin and Milton (1960)
East coast of Mexico	$3.22 \pm 0.03$		Herrin and Richmond (1960)
Czechoslovakia	3.2 Lg		Pec (1961)

Table 11 EARTHQUAKES REVISED

No.	Date			Origin t.			Coordinates		Depth m <sub>b</sub>		Δ	Az	Char.	Lg/P
	m	d	y	h	m	s	Lat	LongW						
Brazil and Peru-Brazil														
1	2	13	64	11	21	44.3	18.0S	56.7	16	5.5	10.9	276	e	5.0
2	6	13	64	3	56	22.4	2.5N	59.9	65	4.5	20.8	204	e	1.0
3	2	22	76	3	24	46.0	0.3N	59.1	10	4.8	19.0	208	e	13.6
4	7	13	76	9	21	45.7	7.4S	73.9	33	4.9	10.7	148	e	2.3
5	8	24	76	0	14	28.3	8.3S	74.4	90	4.7	10.0	148	e	2.4
6	8	2	77	17	45	52.5	0.3S	50.1	33	4.6	24.2	226	e	4.3
7	5	28	78	6	7	3.8	6.7S	74.5	75	4.9	11.6	148	i	3.7
8	3	6	80	9	46	17.7	6.1S	71.2	67	4.8	10.8	164	e	10.8
9	11	12	80	21	23	5.0	8.1S	50.2	33	4.8	19.3	243	e	2.6
10	11	20	80	3	29	41.8	4.5S	38.3	0	5.2	31.6	245	i	1.7
11	6	21	83	9	23	56.2	8.6S	74.4	152	5.1	10.0	143	e	4.0
12	4	8	86	18	2	44.6	7.9S	73.9	173	5.8	10.0	147	e	3.7
13	7	14	86	6	34	44.5	9.6S	72.4	33	4.9	8.0	149	e	10.5
14	11	40	86	5	19	48.3	5.5S	35.7	5	4.9	33.5	248	e	2.8
Colombia														
15	5	12	75	0	0	39	6.9N	73.1	159	4.6	23.8	168	e	1.3
16	6	23	75	5	22	48.3	6.8N	73.1	162	4.9	23.7	152	i	0.6
17	2	1	76	3	3	36.3	0.4N	77.2	42	4.6	19.1	152	e	6.0
18	3	13	76	21	44	41.8	6.8N	73.0	166	5.3	23.7	168	e	2.0
19	5	12	76	16	42	15.1	7.4N	74.9	61	5.1	24.7	164	e	4.4
20	6	14	76	1	37	0.1	6.7N	73.0	161	4.8	23.6	168	e	1.2
21	8	3	76	2	19	22.7	5.3N	75.9	123	4.7	28.8	160	i	1.1
22	1	10	78	20	8	36.1	3.5N	73.6	42	4.8	20.6	165	e	2.5
23	4	28	78	4	28	29.0	12.0N	72.5	13	5.2	28.7	171	e	0.9
24	5	15	78	21	52	43.3	6.2N	77.4	6	4.7	24.4	168	i	0.7
25	5	27	78	16	16	42.6	6.8N	73.0	165	4.7	23.6	178	e	1.5
26	6	18	78	2	20	25.3	6.8N	72.9	169	4.7	23.7	168	i	0.6
27	10	5	78	23	22	21.0	7.4N	76.9	35	4.7	25.3	160	i	0.8
28	5	29	79	12	59	2.5	5.3N	75.7	122	4.9	22.9	161	e	1.1
29	9	2	79	2	0	12.4	4.3N	76.4	101	4.7	22.2	159	i	1.7
30	3	6	80	13	42	9.0	6.0N	74.2	60	4.6	23.2	165	e	1.9
31	5	25	80	15	43	30.4	5.4N	74.5	33	5.0	22.5	164	e	1.9
32	11	18	80	16	34	38.5	6.8N	72.9	171	4.9	23.7	168	i	1.2
33	11	26	80	17	35	41.2	6.9N	72.4	46	4.9	24.6	170	i	3.0
34	4	27	81	18	50	38.7	7.0N	76.6	33	4.9	25.0	160	e	1.4
35	5	13	81	4	38	25.0	4.1N	77.1	47	4.6	22.3	157	i	1.8
36	5	20	81	3	1	43.0	4.7N	76.3	136	4.5	22.6	159	i	1.2
37	6	13	81	18	39	23.9	6.8N	73.0	171	5.0	23.7	168	e	0.5
38	8	5	81	12	58	28.0	3.9N	76.4	62	5.1	21.9	188	e	2.0
39	8	30	81	20	50	9.4	6.9N	76.5	35	4.9	24.7	160	e	1.3
40	10	24	81	4	24	50.9	6.8N	73.0	167	4.6	23.7	168	e	0.8
41	10	26	81	9	5	28.8	6.8N	73.0	165	4.9	23.7	168	i	0.8
42	12	8	81	16	15	22	8.8S	73.1	62	5.1	9.1	148	i	1.6
43	12	17	81	12	54	3.4	6.8N	73.0	165	5.0	23.7	168	e	0.7
44	2	5	82	5	14	36.7	8.1S	74.4	169	4.8	10.3	144	e	0.8
45	2	23	82	20	7	30.8	6.7N	73.0	175	4.7	23.6	168	e	0.8

No.	Date	Origin t.	Coordinates	Depth m <sub>b</sub>	Δ °	Az °	Char.	Lg/P
46	3 9 82	12 21 52.2	6.8N 72.7	170 4.7	23.6	168	e	1.1
47	5 14 82	1 18 50.8	2.4N 75.5	63 4.6	20.1	159	i	4.4
48	7 12 82	13 35 52.0	4.3N 73.5	16 4.6	21.4	166	e	1.7
49	12 23 82	22 47 2.4	6.9N 72.8	160 4.8	23.7	169	e	1.2
50	3 7 83	23 14 11.4	6.9N 73.0	163 4.8	23.8	168	e	1.4
51	3 31 83	13 12 51.0	2.5N 76.7	12 5.4	20.4	156	i	7.2
52	7 24 83	15 39 45.6	6.8N 73.0	165 4.8	23.7	168	e	1.0
53	8 29 83	8 24 24.7	6.8N 73.0	169 5.0	23.7	168	i	1.0
54	12 31 83	12 18 5.5	6.8N 73.0	170 4.7	23.7	168	e	0.7
55	1 6 84	11 37 49.8	6.7N 73.1	166 5.0	23.6	168	e	1.3
56	1 25 84	18 46 25.0	3.5N 76.7	49 4.6	31.6	157	e	1.2
57	1 28 84	17 4 39.2	6.7N 74.5	81 5.0	23.9	165	e	1.6
58	8 11 84	13 14 20.9	6.8N 73.0	173 4.7	23.6	168	e	0.6
59	10 27 84	11 56 13.2	9.8N 74.7	61 5.0	27.0	166	e	1.1

Venezuela

60	6 12 74	16 25 45.2	10.6N 63.5	11 5.7	27.3	190	e	1.4
61	9 20 74	14 47 57.9	9.3N 70.6	41 4.7	25.8	175	e	2.2
62	10 29 74	3 10 16.9	10.6N 63.4	33 5.0	27.1	191	e	2.5
63	3 5 75	13 47 58.3	9.1N 69.9	25 5.5	25.5	176	Disturbed	
64	4 5 75	9 34 37.6	10.1N 68.9	36 5.5	26.2	177	e	8.0
65	7 18 75	7 17 33	10.9N 64.5	3 4.8	27.3	188	e	1.2
66	8 24 75	1 5 15.1	10.7N 62.6	111 5.1	27.4	191	e	1.1
67	12 5 75	9 31 50.8	10.8N 62.7	114 4.9	27.5	191	i	1.6
68	12 2 76	5 33 54.3	10.8N 63.7	38 4.8	26.9	177	i	2.0
69	12 21 76	4 32 31	8.8N 61.7	40 4.7	25.7	194	e	2.9
70	2 21 77	3 7 43.5	10.5N 62.5	45 4.7	27.4	192	e	1.1
71	2 21 77	17 48 1.0	9.5N 70.8	4 5.0	26.0	174	e	3.0
72	7 24 77	5 44 44.3	10.8N 68.8	14 4.6	26.9	179	i	3.2
73	8 14 77	4 22 49.7	10.9N 62.4	110 4.2	27.6	192	e	2.0
74	9 3 77	15 25 16.1	10.4N 62.3	35 4.7	27.1	192	e	3.2
75	9 14 77	20 51 8.8	10.8N 62.4	94 4.7	27.5	192	e	1.3
76	9 18 77	17 31 16.2	10.5N 63.3	42 4.6	27.0	190	e	2.2
77	12 11 77	16 22 6.2	9.6N 69.5	2 5.5	25.7	177	i	4.5
78	12 17 77	23 25 10	10.9N 65.5	14 4.6	27.4	186	i	4.0
79	1 18 78	1 17 54.5	10.3N 62.2	46 4.8	27.3	192	i	1.1
80	3 15 78	15 26 37.0	10.3N 62.2	11 4.5	27.3	192	e	6.2
81	5 18 78	3 25 04.9	10.8N 62.5	116 4.7	27.7	192	e	1.2
82	11 7 78	2 40 23	8.6N 62.9	17 4.6	25.5	192	c	2.8
83	3 30 79	12 10 7.0	12.9N 70.8	33 4.6	29.4	175	e	1.4
84	5 5 79	20 4 56.0	8.4N 70.9	8 5.4	25.0	174	e	3.7
85	5 5 79	20 8 40.3	8.5N 70.5	34 5.2	25.0	173	e	5.0
86	7 17 79	8 49 28.8	10.2N 62.2	40 4.6	27.0	193	e	1.4
87	8 3 79	11 43 57.3	8.7N 70.8	15 4.8	25.0	174	e	1.4
88	2 12 80	2 29 14.0	9.8N 68.6	24 4.6	26.2	179	e	2.3
89	11 17 80	16 50 21.5	10.9N 69.5	39 4.6	27.3	177	i	2.0
90	12 20 80	17 0 24.3	9.7N 72.4	66 4.6	26.4	171	e	1.1
91	6 23 81	22 57 39.0	10.5N 63.4	11 5.0	27.0	190	e	2.3
92	10 18 81	4 31 1.2	8.2N 72.5	37 5.5	24.7	170	e	3.3

No.	Date	Origin t.	Coordinates	Depth m <sub>b</sub>	Δ °	Az°Char.	Lg/P
93	12 25 81	12 35 48.3	10.9N 62.4	96 5.1	27.9	192 e	1.8
94	1 15 82	3 59 18.0	9.4N 69.9	12 5.1	25.6	176 e	2.0
95	3 18 82	2 11 50.0	10.5N 62.4	58 4.7	27.4	192 e	1.6
96	5 10 82	1 25 57.3	10.9N 62.5	100 5.2	27.4	192 no	
97	5 27 82	11 26 6.6	8.7N 70.3	14 4.7	25.0	174 e	1.2
98	8 10 82	8 24 0	10.7N 62.6	104 4.8	27.3	192 e	1.3
99	11 23 82	17 27 1.0	10.6N 63.2	23 4.8	27.4	190 e	1.2
100	12 11 82	10 18 37.3	8.6N 71.7	14 5.1	25.3	172 e	1.4
101	3 19 83	3 0 26.3	10.6N 63.2	28 4.6	27.4	190 i	1.6
102	4 11 83	8 6 7.2	10.5N 62.7	43 4.7	27.3	191 i	4.0
103	4 11 83	8 18 10.2	10.4N 62.7	38 5.9	27.3	191 e	1.8
104	5 2 83	21 55 52.4	10.3N 62.3	45 4.5	27.0	192 e	1.6
105	1 23 84	21 36 50.8	10.7N 62.7	120 5.4	27.3	191 e	1.7
106	5 25 84	0 59 23.6	10.4N 62.4	41 4.7	27.1	192 i	3.0
107	6 12 84	23 8 55.5	7.9N 71.3	38 4.7	24.2	173 e	0.9
108	6 14 84	10 4 30.5	9.9N 69.8	38 5.2	26.1	176 e	10.7

Chile

109	5 15 72	9 12 56.6	29.7S 71.3	49 4.9	13.4	13 e	0.8
110	5 28 72	7 28 13.5	27.7S 71.3	41 4.8	11.5	15 e	0.2
111	5 28 72	9 46 15.4	27.7S 71.4	4 4.9	11.5	16 e	2.3
112	10 26 82	3 24 30.1	29.7S 71.4	63 5.6	13.4	14 e	1.8
113	1 26 86	7 48 23.5	27.0S 70.9	18 5.7	10.8	14 e	3.3
114	3 21 86	13 55 41.0	30.7S 71.4	52 4.9	17.0	13 e	0.7
115	5 14 86	15 54 23.8	32.6S 71.9	33 5.0	17.0	13 e	0.6
116	5 19 86	12 36 30.3	28.4S 69.1	113 5.1	15.0	5 e	0.8
117	6 24 86	12 25 28.3	30.7S 71.7	51 5.4	17.0	14 e	0.6
118	7 28 86	3 29 56.0	33.3S 72.0	41 4.7	19.0	13 e	0.8
119	7 28 86	20 29 2.7	33.3S 71.9	41 5.1	19.0	13 e	0.5

Argentina

120	1 7 74	16 35 56.0	26.9S 65.7	20 5.7	10.5	347 e	0.8
121	8 16 74	7 47 51.5	33.3S 68.3	35 4.8	16.7	359 e	0.8
122	8 24 74	18 58 20.0	31.4S 67.4	12 5.3	14.8	357 e	0.8
123	9 3 74	20 22 20.5	25.9S 67.6	45 4.8	9.3	357 e	0.8
124	10 6 74	14 40 56.1	30.9S 65.1	40 4.7	14.6	348 e	0.8
125	5 6 75	18 10 2.0	32.9S 69.0	14 5.0	16.3	357 e	0.8
126	1 4 76	4 42 4.0	27.9S 66.0	128 4.8	11.5	350 e	0.8
127	2 14 76	9 1 52	33.6S 68.9	20 4.8	17.1	357 e	0.8
128	3 20 76	2 55 47.8	27.6S 67.4	118 4.8	11.0	356 e	0.8
129	3 27 76	21 5 7.1	31.8S 67.7	122 5.1	15.2	358 e	0.8
130	5 4 76	2 7 11.3	27.3S 65.8	58 4.7	11.0	348 e	0.8
131	8 3 76	23 43 54.6	31.5S 68.5	119 5.0	14.9	358 e	0.8
132	10 24 76	0 13 51	32.8S 69.2	25 4.9	16.2	356 e	0.8
133	11 26 76	7 13 38	28.0S 64.7	25 5.0	11.8	344 e	0.8
134	11 23 77	11 8 43	31.3S 67.8	32 5.3	14.7	359 e	0.8
135	11 23 77	11 44 23	31.6S 67.8	60 4.7	15.0	359 Disturbed	
136	11 23 77	11 46 55.4	31.0S 67.6	8 5.2	14.4	359 e	0.9
137	11 23 77	11 58 10.0	31.0S 67.9	22 5.5	14.4	359 e	0.9

No.	Date	Origin t.	Coordinates	Depth m <sub>b</sub>	Δ	Az°	Char.	Lg/P
138	11 23 77	13 58 51	31.5S 68.2	49 4.8	14.9	360	e	0.3
139	11 23 77	15 37 56	31.8S 67.7	65 4.6	15.2	359	e	0.3
140	11 23 77	16 17 51	31.2S 67.5	20 4.5	14.6	358	e	0.2
141	11 23 77	16 28 23	31.3S 67.7	10 5.1	14.7	358	e	0.3
142	11 23 77	16 36 3.0	31.3S 67.7	21 5.6	14.7	359	e	0.8
143	11 23 77	21 52 2.5	31.3S 67.7	36 4.9	14.8	359	no	
144	11 23 77	21 57 28.3	31.7S 67.7	62 4.9	15.1	359	e	0.4
145	11 23 77	23 4 13.4	31.8S 68.0	84 5.0	15.2	359	e	0.5
146	11 23 77	23 27 36.0	31.6S 67.7	20 5.1	15.0	358	e	0.8
147	11 24 77	1 57 32.2	31.6S 67.7	19 5.5	15.0	358	Disturbed	
148	11 24 77	3 20 5.0	31.7S 67.8	62 4.4	15.1	359	e	0.4
149	11 24 77	3 57 50.0	31.8S 68.8	48 4.3	15.2	359		0.3
150	11 24 77	5 6 24.9	31.7S 67.6	45 4.3	15.2	358	e	0.4
151	11 24 77	6 13 35.0	31.6S 67.8	19 4.2	15.0	358	e	0.5
152	11 24 77	11 8 39	31.7S 67.9	83 4.0	15.1	359	e	0.3
153	11 24 77	18 19 14.6	31.4S 67.6	33 4.8	14.8	358	Disturbed	
154	11 24 77	18 20 16.5	31.3S 67.7	33 5.6	14.7	358	e	0.8
155	11 24 77	18 42 40	31.4S 67.8	26 5.8	14.8	359	e	0.8
156	11 24 77	22 19 58.3	31.5S 67.7	51 4.4	14.9	359	Disturbed	
157	11 24 77	23 0 54.5	31.2S 67.8	47 4.9	14.6	359	e	0.8
158	11 25 77	0 4 31.6	31.1S 67.7	43 5.4	14.5	359	Disturbed	
159	11 25 77	3 24 37.3	31.7S 67.7	33 4.6	15.1	359	e	0.8
160	11 25 77	3 47 16.6	31.8S 67.6	41 5.0	15.2	358	e	0.8
161	11 25 77	4 6 53.1	31.2S 67.7	22 4.8	14.6	358	e	0.8
162	11 25 77	18 2 40.3	31.2S 67.8	53 4.9	14.6	359	e	0.8
163	11 25 77	18 56 32.1	31.4S 67.5	33 4.9	14.8	358	e	0.8
164	11 25 77	20 42 15.6	31.3S 67.6	47 4.1	14.7	358	e	0.4
165	11 26 77	13 52 30.6	31.3S 67.5	33 5.0	14.7	358	e	0.3
166	11 26 77	20 26 51.0	31.3S 67.7	33 3.7	14.8	358	e	0.3
167	11 27 77	6 26 4.0	31.1S 67.7	33 3.6	14.5	358	no	
168	11 27 77	10 15 5.3	31.6S 67.8	47 4.8	15.0	359	no	
169	11 28 77	0 17 24.3	31.0S 67.7	28 5.3	14.4	358	no	
170	11 28 77	4 19 31.0	31.7S 67.6	2 5.6	15.1	358	no	
171	11 28 77	5 39 24.0	31.0S 68.0	23 5.3	14.4	360	no	
172	11 28 77	6 31 29.1	31.4S 67.4	17 5.9	14.8	358	no	
173	11 28 77	18 40 18.8	31.9S 69.0	97 5.2	15.3	358	no	
174	11 28 77	23 7 57	31.7S 67.3	1 4.9	15.1	357	e	0.4
175	11 29 77	0 33 38	31.6S 67.8	33 4.8	15.0	359	e	0.4
176	12 5 77	15 43 26.0	31.1S 68.0	11 5.4	14.5	359	e	0.8
177	12 6 77	8 41 35	31.0S 67.7	5 5.4	14.4	359	e	0.6
178	12 6 77	12 41 16.8	31.4S 67.8	33 4.9	14.8	359	no	
179	12 6 77	17 5 6.9	31.2S 67.9	21 5.9	14.6	359	e	0.5
180	12 6 77	18 27 38	31.3S 67.6	4 5.1	14.7	358	e	0.3
181	12 7 77	3 22 44	31.2S 67.9	27 5.1	14.6	359	e	0.8
182	12 10 77	7 11 55.6	31.3S 67.7	39 5.1	14.7	358	e	0.4
183	12 10 77	8 37 1.0	31.3S 67.7	37 4.8	14.7	359	e	0.4
184	12 10 77	14 19 57.9	31.2S 67.7	27 5.2	14.6	359	e	0.7
185	12 12 77	16 2 33	31.4S 67.5	40 4.9	14.3	358	e	0.6
186	12 21 77	3 47 32.2	31.6S 67.7	33 5.7	15.0	359	e	0.6
187	1 1 78	10 50 56	31.1S 67.9	17 5.1	14.5	359	e	0.6



No.	Date	Origin t.	Coordintes	Depth m <sub>b</sub>	Δ °	Az °	Char.	Lg/P
188	1 3 78	1 10 4.4	31.5S 67.9	35 5.2	14.9	359	e	0.6
189	1 3 78	6 31 5.1	31.3S 67.9	38 5.0	14.7	359	e	0.6
190	1 17 78	11 33 14.5	31.2S 68.0	20 5.8	14.7	360	e	0.6
191	1 22 78	12 8 26.8	31.5S 67.9	35 4.6	14.8	359	e	0.3
192	1 24 78	13 18 15.7	31.7S 68.9	18 5.6	15.1	357	e	0.6
193	3 18 78	0 38 40	31.4S 67.8	16 5.1	14.2	359	e	0.6
194	4 4 78	19 33 53.3	31.2S 67.7	17 5.4	14.6	359	e	0.6
195	5 10 78	23 6 2.0	30.0S 68.9	38 5.1	13.4	357	e	0.8
196	6 7 78	15 16 45.0	32.1S 67.6	44 5.1	15.5	358	e	0.4
197	6 26 78	18 49 11.8	31.6S 67.7	0 5.1	15.0	359	e	0.8
198	7 26 78	1 47 16.1	31.5S 67.8	46 5.0	14.9	359	e	0.9
199	8 21 78	0 28 25.1	31.3S 67.9	25 5.5	14.7	359	e	0.5
200	10 25 78	1 36 41.7	31.5S 67.7	43 5.3	14.9	358	e	0.6
201	1 29 79	3 22 45.5	31.3S 68.4	10 5.0	14.7	359	e	0.4
202	7 11 79	23 8 3.1	31.5S 78.3	51 4.8	14.6	359	e	0.6
203	8 30 79	18 59 46.9	31.5S 67.7	47 5.4	14.9	358	e	0.8
204	10 8 79	1 52 40.0	31.5S 68.0	39 4.8	14.9	360	e	0.8
205	1 17 80	11 0 10.0	31.5S 67.7	29 4.8	14.9	359	e	0.8
206	1 24 80	18 34 4.1	31.8S 68.5	43 4.7	15.2	359	e	0.8
207	4 9 80	8 17 57.4	31.6S 67.5	23 5.4	15.1	358	e	0.7
208	5 25 80	21 46 11.8	31.3S 68.0	43 5.0	14.7	360	e	0.7
209	11 10 80	16 24 39.0	31.6S 67.5	13 5.6	15.0	358	e	0.7
210	12 6 80	3 43 11.5	31.3S 67.5	46 4.8	14.7	358	e	0.7
211	7 2 81	11 3 35	33.0S 69.1	58 4.5	16.4	357	e	0.7
212	8 4 82	5 12 20.2	30.5S 68.1	53 4.8	13.9	360	e	0.6
213	12 4 82	3 26 42.6	31.3S 67.7	36 4.9	14.7	359	e	0.6
214	5 6 84	13 57 31.7	30.6S 68.9	81 4.6	14.0	357	e	0.6

Perú

215	8 12 82	8 27 28.0	6.7S 75.8	33 4.7	12.3	143	i	3.1
216	8 15 82	6 11 16.8	10.1S 76.5	117 5.5	10.3	129	e	4.0
217	2 27 83	5 5 19.0	13.5S 76.8	33 5.4	8.9	111	e	0.6
218	3 20 83	1 56 38.6	10.5S 74.9	18 5.4	8.9	133	e	1.5
219	3 21 83	1 2 51.0	3.7S 78.3	46 4.5	16.2	143	e	1.5
220	3 9 86	16 47 51.8	8.1S 80.1	33 4.9	15.3	127	no	
221	4 2 86	6 49 30.4	4.1S 80.8	33 4.7	18.0	135	no	
222	4 23 86	20 27 12.7	3.9S 80.9	46 5.4	18.0	136	e	0.5
223	4 28 86	13 43 12.0	15.0S 75.5	57 4.8	8.1	103	e	0.6
224	5 21 86	3 56 56.2	3.4S 76.7	121 4.9	18.0	148	e	1.2
225	7 17 86	18 11 22.9	9.2S 79.9	33 4.9	14.0	124	e	1.6
226	7 20 86	5 52 16.2	15.9S 75.3	33 4.6	9.1	96	no	
227	7 21 86	0 40 46.8	14.1S 76.2	43 4.1	8.0	108	no	
228	7 27 86	2 54 34.4	10.0S 76.3	33 4.7	9.0	130	no	

Table III Lg MEAN APPARENT VELOCITY

<u>Origen Zone</u>	<u>Velocity (Km/sec)</u>
Brazil	$3.60 \pm 0.2$
Colombia	$3.60 \pm 0.2$
Venezuela	$3.58 \pm 0.2$
Chile	$3.68 \pm 0.3$
Argentina	$3.52 \pm 0.2$
Peru	$3.60 \pm 0.3$




Figure 1: Simplified Geological map of South America. (Martinez, 1980; Almeida and Hasui, 1984)




Figure 2: Map of "Lg" paths across South America to LPB station

COMP 2 SP A = 50 K

BRAZIL 02/22/76

Lg 03h. 34min. 42sec.

COMP 2 SP A = 25 K

BRAZIL 05/28/78

Lg 06h. 13min. 13sec.

COMP 2 SP A = 50 K

COLOMBIA 03/13/76

Lg 21h. 56min. 36sec.

Fig. 3 Earthquakes recorded at LPB station, travel across the shield.





Fig. 4 Earthquakes recorded at LPB station, travel across the shield.

COMP  $\pm$  SP  $\Delta$  = 50 K

CHILE 05/28/72

Lg 07h. 34min. 20sec.

COMP  $\pm$  SF  $\Delta$  = 25 K

ARGENTINA 08/04/82

Lg 05h. 20min. 00sec.

COMP  $\pm$  SP  $\Delta$  = 25 K

PERU 08/12/82

Lg 08h. 34min. 02sec.

Fig. 5 Earthquakes recorded at LPB station, travel across the Andes.

# PARTICLE MOTION





Figure 6: Horizontal and vertical components particle motion diagram:  
a) Brazil earthquake 11/20/80  
b) Colombia earthquake 5/11/82



# PARTICLE MOTION

a)



b)





Fig. 7 Horizontal and vertical components particle motion diagrams:

a) Venezuela earthquake 7/17/79

b) Chile earthquake 6/28/72

# PARTICLE MOTION

a)



b)




Figure 8: Horizontal and vertical components particle motion diagrams:

a) Argentina earthquake 12/05/77

b) Peru earthquake 6/27/83

# SPECTRUM

-30-





Figure 9: Lg spectra for earthquakes originating in Brazil and Colombia





Fig. 10 Lg spectra for earthquakes originated in Venezuela and Chile





Fig. 11 Lg spectra for earthquakes originated in Argentina and Peru